

Pouchoscopy Prep (with Fleet Enema)

The following instructions will help you prepare for your pouchoscopy. **Please read all instructions in this packet at least 1 week prior to your procedure.**

Date:

Procedure time:

Arrival time:

Physician:

Location:

Arrival

- **Do not go to your doctor's office.**
- **Newton-Wellesley Hospital:** Enter through the Surgical Center/East Entrance, proceed to GI registration desk.
- **Boston Endoscopy Center:** Do not go to Newton-Wellesley Hospital. Go to 175 Worcester St, Wellesley Hills.
- **St. Elizabeth's Medical Center:** Present to the Admissions Office for registration.
- Please note that procedure times are not exact and unexpected delays may occur.

Transportation

- This procedure may be done with or without sedation.
- If you choose to have sedation, you must arrange a ride home before the day of your procedure. After your exam, you will not be able to drive or take any other transportation by yourself. You must arrange for a responsible adult to accompany you by car, taxi or public transportation. This is strictly observed for your safety. No exceptions will be made unless you have the procedure without sedation.

Cancellation/No Show Policy for Procedures

- Scheduling time for procedures requires planning and coordination between the staff, doctors, and the hospital. Since the schedule is often carefully planned to avoid delays and having patients taken care of in a timely fashion, it is important that you adhere to the schedule as arranged. Should you need to reschedule or cancel, please do so at least 7 days prior to procedure.
- If you have your menstrual period, the exam will not be affected. However, if you have a fever, please call the office to reschedule.
- If a procedure is not cancelled **at least 7 days in advance**, you will be charged a one-hundred-dollar (\$100) fee. This fee is NOT covered by your insurance company.

Medication Concerns

- If you use **antiplatelets or anticoagulants** such as apixaban (Eliquis), clopidogrel (Plavix), dabigatran (Pradaxa), dipyridamole (Persantine), edoxaban (Savaysa), prasugrel (Effient), rivaroxaban (Xarelto), ticlopidine (Ticlid), or warfarin (Coumadin), please notify us immediately.
- Stop taking **iron supplements** and **fiber supplements** 5 days prior to procedure.
- Continue taking daily aspirin if prescribed.

Preparation Instructions

You will need the following:

- One fleet enema

1 day before procedure (before 5 p.m.)

- You may have a normal breakfast and lunch.

Evening before procedure (between 5–9 p.m.)

- Begin a clear liquid diet. Clear liquids include water, fruit juices without pulp (apple or white grape juice), carbonated drinks (ginger ale, seltzer, sodas), gelatin, tea or coffee (without milk or cream), sports drinks, fat-free broth, bouillon, frozen juice popsicles.
- Avoid red or purple liquids.

Morning of procedure

- Have clear liquids only for breakfast.
- Avoid red or purple liquids.
- Administer one fleet enema at least 2 hours before your scheduled procedure time.
- If you are having this procedure done with sedation, **finish clear liquids at least 3 hours before procedure.**